

# IMMUNISATION SCHEDULE

Sprache: Englisch

The vaccinations should be administered at the earliest point in time possible.

Checking the vaccination status makes sense at any age. Missing vaccinations should be caught up on immediately according to the recommendations for the respective age.

Recommendations of the Standing Committee on Vaccination (STIKO), 2024. [www.stiko.de](http://www.stiko.de)



| VACCINATION                                | AGE | in weeks        |    | in months |   |                 |     |                 |       |    | in years |     |                                 |     |      |       |                |                                 |                 |                |                |  |
|--|-----|-----------------|----|-----------|---|-----------------|-----|-----------------|-------|----|----------|-----|---------------------------------|-----|------|-------|----------------|---------------------------------|-----------------|----------------|----------------|--|
|  |     | 6               | 2  | 3         | 4 | 5-10            | 11* | 12              | 13-14 | 15 | 16-23    | 2-4 | 5-6                             | 7-8 | 9-14 | 15-16 | 17             | from 18                         | from 60         |                |                |  |
|  |     | U4              |    | U5        |   |                 | U6  |                 |       |    | U7       |     | U7a/U8                          |     | U9   |       | U10            |                                 | U11/J1          |                | J2             |  |
| Rotavirus                                  |     | G1 <sup>a</sup> | G2 | (G3)      |   |                 |     |                 |       |    |          |     |                                 |     |      |       |                |                                 |                 |                |                |  |
| Tetanus <sup>b</sup>                       |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    | N        |     |                                 |     | A1   | N     | A2             | N                               |                 |                | A <sup>f</sup> |  |
| Diphtheria <sup>b</sup>                    |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    | N        |     |                                 |     | A1   | N     | A2             | N                               |                 |                | A <sup>f</sup> |  |
| Pertussis <sup>b</sup>                     |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    | N        |     |                                 |     | A1   | N     | A2             | N                               | A3 <sup>f</sup> | N              |                |  |
| Haemophilus influenzae Type b <sup>b</sup> |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    | N        |     |                                 |     |      |       |                |                                 |                 |                |                |  |
| Poliomyelitis <sup>b</sup>                 |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    |          |     |                                 | N   |      |       |                | A1                              |                 |                | N              |  |
| Hepatitis B <sup>b</sup>                   |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    |          |     |                                 |     |      |       | N              |                                 |                 |                |                |  |
| Pneumococci <sup>b</sup>                   |     | G1              | N  | G2        | N | G3 <sup>c</sup> |     |                 |       |    | N        |     |                                 |     |      |       |                |                                 | S <sup>h</sup>  |                |                |  |
| Meningococcus B <sup>d</sup>               |     | G1              | N  | G2        | N |                 |     | G3 <sup>d</sup> |       |    |          |     | N                               |     |      |       |                |                                 |                 |                |                |  |
| Measles                                    |     |                 |    |           |   |                 | G1  | N               | G2    |    |          |     |                                 |     |      |       | N              |                                 |                 | S <sup>g</sup> |                |  |
| Mumps, rubella                             |     |                 |    |           |   |                 | G1  | N               | G2    |    |          |     |                                 |     |      |       | N              |                                 |                 |                |                |  |
| Varicella                                  |     |                 |    |           |   |                 | G1  | N               | G2    |    |          |     |                                 |     |      |       | N              |                                 |                 |                |                |  |
| Meningococcal C                            |     |                 |    |           |   |                 |     | G1              |       |    |          |     |                                 |     |      |       |                |                                 | N               |                |                |  |
| Human papillomavirus                       |     |                 |    |           |   |                 |     |                 |       |    |          |     | G1 <sup>e</sup> G2 <sup>e</sup> |     |      | N     |                |                                 |                 |                |                |  |
| COVID 19                                   |     |                 |    |           |   |                 |     |                 |       |    |          |     |                                 |     |      |       | G <sup>i</sup> | S <sup>i</sup>                  |                 |                |                |  |
| Herpes zoster                              |     |                 |    |           |   |                 |     |                 |       |    |          |     |                                 |     |      |       |                | G1 <sup>j</sup> G2 <sup>j</sup> |                 |                |                |  |
| Influenza                                  |     |                 |    |           |   |                 |     |                 |       |    |          |     |                                 |     |      |       |                | S (annually)                    |                 |                |                |  |

Next vaccination appointment



## EXPLANATIONS

- G** BASIC IMMUNISATION  
(in up to 3 inoculations G1 – G3)
- S** STANDARD VACCINATION
- A** BOOSTER VACCINATION
- N** CATCH-UP VACCINATION  
(Basic immunisation of all persons not yet vaccinated or completion of an incomplete series of vaccinations)
- U** Early detection examination
- J** Youth examination  
(J1 at the age of 12–14 years old)

<sup>a</sup> First vaccine dose as early as the age of 6 weeks; depending on the vaccine used 2 or 3 vaccine doses at an interval of at least 4 weeks.  
<sup>b</sup> Premature infants: Additional vaccine dose at the age of 3 months, i.e. a total of 4 vaccine doses.  
<sup>c</sup> Minimum distance from the previous dose: 6 months.  
<sup>d</sup> 3 doses at the age of 2 to 23 months; from the age of 24 months, the vaccination series consists of 2 doses.  
<sup>e</sup> Two vaccine doses at least 5 months apart; in the case of a catch-up immunisation starting at an age ≥ 15 years old or a vaccination interval of < 5 months between the 1st and 2nd dose, a 3rd dose is necessary.

<sup>f</sup> Td booster vaccination every 10 years. Next due Td vaccination requires one dose as a Tdap or, in the case of a respective indication, as a Tdap-IPV combination vaccination.  
<sup>g</sup> One-time vaccination with an MMR vaccine for all persons born after 1970 ≥ 18 years old with an ambiguous vaccination status, without prior vaccination or with only one vaccination during childhood.  
<sup>h</sup> Recommended with PCV20  
<sup>i</sup> For the protection of individuals without an underlying disease, (at least) 3 antigen contacts are required, at least one of which must be a vaccination.

<sup>j</sup> Repeated booster vaccination with a minimum interval of 12 months from the last antigen contact, preferably in autumn  
 \* Vaccinations can be spread over multiple vaccination appointments. MMR and V can be administered at the same appointment or in 4-week intervals.